



91998000385900A703200000000050307PCTKR2004002255

[별지 제65호의48 서식]

P C T

방식 심사 사관	담 당	심 사 관

【Title of Document】 SUBMISSION OF AMENDMENTS

【Receiver】 Commissioner of the Korean Intellectual Property Office

【Identification of International Application】

【International Application No.】 PCT/KR2004/002255

【International Filing Date】 2004.09.06

【Priority Date】 2003.09.06.

【Applicant】

【Name】 OSCOTEC INC. et al.

【Address】 2-17, Omok-ri, Seonggeo-eup, Cheonan-si,

Chungcheongnam-do 330-831, Republic of Korea

【Tel. No.】 82-2-3453-0507

【Applicant 's code】 1-1999-036103-0

【State of Nationality】 KR

【Agent】

【Name】 LEE, Won-Hee

【Address】 8th Fl., Sung-ji Heights II, 642-16 Yoksam-dong,

Kangnam-ku, 135-080, Seoul, Republic of Korea

【Tel. No.】 82-2-3453-0507

【Fax No.】 82-2-3453-8155

【Agent 's Code】 9-1998-000385-9

【Purport】 Submitted hereby is a correction pursuant to Article

106-36(3) of the Enforcement Regulations of the Patent Law.

Agent

LEE, Won-Hee (Seal)

【Attached Document(s)】 1.replacement sheet for correction_2Copies



LETTER FOR PCT ARTICLE 34 AMENDMENT

Date : 2005. 8. 3

Amendment of the claims under Article 34

International Application No. : PCT/KR2004/002255

International Filing Date : 6 September 2004

Applicant : OSCOTEC INC. et al.

Agent : LEE, Won-Hee

8th Fl., Sung-ji Heights II 642-16, Yoksam-dong,

Kangnam-ku, 135-080, Seoul, Republic of Korea

Applicant's or Agent's File reference : 4fpo-08-02

Dear Sir(s) :

Applicant herewith submits replacement sheets numbered pages 10-39 to replace sheets numbered pages 10-38 originally filed for this application.

In respect of each claim appearing in the international application based on the replacement sheets submitted herewith, and in accordance with PCT Section 205, the following claim(s) is/are :

(i) amended : Claims 1 to 4

In claim 1, "*Notoginseng radix* extract" is changed into "a purified form of butanol fraction of ethanol extract of *Notoginseng radix*", the ground of which can be found in <1-2> Separation of *Notoginseng radix* crude extract in page 19 of original specification.

In claim 2, 'product by process' claim has been created.

In claim 3 and 4, amended claims are just renumbering of original claim 4 and 5.

(ii) deleted : Claim 5

Sincerely yours,

Won-hee, Lee (Seal)

oil composition is fewer in *Notoginseng radix* than in *Panax ginseng*. *Notoginseng radix* additionally includes oleanolic acid. Its root has hemostatic and cardiotonic activities. It was confirmed from animal tests that the root has efficacy of increasing blood flow of coronary artery, decreasing oxygen consumption of cardiac muscle and lowering the levels of lipid and cholesterol in blood. *Notoginseng radix* also has functions of anti-inflammation, analgesia and hemostasis, so that it is very useful for the treatment of not only inflammatory diseases including hepatitis but also bleeding from trauma, cut, etc., and internal hemorrhage. Applying to a wound or oral administration give the same effects.

Notoginseng radix extract of the present invention is extracted by using water, alcohol or a mixed solvent of water and alcohol. At this time, alcohol is preferred to be ethanol.

Conventional extraction methods including cold precipitation, hot precipitation, heating, etc, using the solvent mentioned above are used.

Notoginseng radix extract of the present invention is more preferably a purified form of butanol fraction of ethanol extract of *Notoginseng radix*.

The purified form of butanol fraction of ethanol extract of *Notoginseng radix* may be produced by the steps of: obtaining ethanol extract of *Notoginseng radix* by using ethanol as an extraction solvent;
5 obtaining butanol fraction of ethanol extract from the ethanol extract by using butanol as fractionation solvent; separating an intermediate form of butanol fraction of ethanol extract from the butanol fraction by using column chromatography; and separating a
10 purified form of butanol fraction of ethanol extract from the intermediate form of butanol fraction by using column chromatography.

Notoginseng radix extract of the present invention inhibits release of tumor necrosis factor-alpha (TNF- α), so that it can be used for the
15 production of health food or a medicine for preventing and treating arthritis.

In order to investigate how *Notoginseng radix* extract of the present invention worked to inhibit
20 release of tumor necrosis factor-alpha (TNF- α), THP-1 cells, a human monocytic cell line, were treated with lipopolysaccharide (LPS) and *Notoginseng radix* extract of the present invention at the concentration of 2 or 10 $\mu\text{l/ml}$. Then, the amount of released tumor necrosis
25 factor-alpha (TNF- α) in cell culture medium was

measured by ELISA. As a result, the amount of released tumor necrosis factor-alpha (TNF- α) was remarkably decreased by the treatment of 10 $\mu\text{l/ml}$ of *Notoginseng radix* extract of the present invention (see
5 Experimental Example 1).

Notoginseng radix extract of the present invention can be used for the production of health food or a medicine for preventing and treating arthritis owing to its ability to death activated T-cells
10 selectively.

In order to investigate whether or not *Notoginseng radix* extract of the present invention was able to death activated T-cells only, a lymph node of a 5-week-old female mouse was taken and single cells were
15 prepared. The cells were cultured, during which T cells were activated. The apoptosis of activated T-lymphocytes was investigated. As a result, when cells were treated with over 5 $\mu\text{l/ml}$ of *Notoginseng radix* extract of the present invention, only activated T-
20 cells were killed, (inactivated T-cells were still alive) (see Experimental Example 2).

Notoginseng radix extract of the present invention also inhibits the progress of the disease in animals having type 2 collagen induced arthritis.

In order to investigate the treatment effect on arthritis of *Notoginseng radix* extract of the present invention, collagen suspension was intra-dermally injected in tail head of a mouse to induce arthritis.

5 *Notoginseng radix* extract of the present invention was orally administered to the mouse with arthritis, which was then observed. As a result, the progress of arthritis was remarkably inhibited from the 9th day after oral administration of the extract (see

10 Experimental Example 3).

A composition of the present invention can additionally include, in addition to *Notoginseng radix* extract, one or more effective ingredients having a similar to or the same function as *Notoginseng radix*

15 extract.

A composition of the present invention can additionally include, in addition to *Notoginseng radix* extract, one or more effective ingredients having a different function from that of *Notoginseng radix*

20 extract.

A composition of the present invention can contain at least one of pharmaceutically acceptable carriers, in addition to the above effective ingredients, for the convenience of the administration.

25 Pharmaceutically acceptable carriers can be selected

from a group consisting of saline, sterile water, Ringer's solution, buffered saline, dextrose solution, maltodextrin solution, glycerol, ethanol and a mixture of them (one or more components). If necessary, other
5 additives such as anti-oxidants, buffers, fungistats, etc, can be included. A composition of the present invention can also be prepared in the forms of pills, capsules, granules, tablets and injectable solutions such as aqueous solutions, suspensions, emulsions, etc,
10 produced by being mixed with generally used diluents, disintegrating agents, surfactants, binders and lubricants. Besides, a composition of the present invention can be prepared in different forms considering a disease and included ingredients by
15 general method well-known to the people in this field or the method described in Remington's Pharmaceutical science (Newest edition), Mack Publishing Company, Easton PA. Calcium or vitamin D₃ can be added to a composition of the present invention to enhance its
20 medicinal effect of preventing and treating arthritis.

The administration method of a composition of the present invention varies from the purpose of the treatment; either oral administration or parenteral administration (for example, intravenous, intradermal,
25 intraperitoneal or local injection) is fine. And the

dosage of the composition is determined according to weight, age, gender, health condition of a patient, diet, administration times and method, excretion rate, and severity of a disease. The effective dosage of
5 *Notoginseng radix* extract of the present invention is 0.1~10 mg/kg, and 0.1~3 mg/kg is more preferable. The administration times can be once a day or preferably several times a day.

The acute toxicity test in mice via oral
10 administration was performed to see if the *Notoginseng radix* extract of the present invention has acute toxicity in mice. As a result, its estimated LD₅₀ values are much greater than 2 g/kg in mice, indicating that this extract is evaluated to be a safe substance.

15 A composition of the present invention can be treated for preventing and treating arthritis either independently or in combination with surgical operation, radiotherapy, hormone therapy, chemotherapy and other biological response regulators.

20 A composition of the present invention can be added to health food to improve arthritis related diseases. *Notoginseng radix* extract of the present invention can be added to food as it is or together with other food or food ingredients by general method
25 for food process. The mixing ratio of effective

ingredients is determined by the purpose of use (for prevention, for promoting health, or for treatment of a disease). In general, *Notoginseng radix* extract of the present invention is added to food or beverages under 100 weight%, preferably under 50 weight%. However, in the case of long-term administration for the purpose of health and sanitation or health control, the amount of a composition added to food or beverages might be less than the above, but since the composition is safe for human, it could be added more than the above.

There is no limitation in food category applicable to the extract of the present invention. So, the extract can be added to meat, sausages, bread, chocolate, candies, snacks, cookies, pizza, ramyun, noodles, gums, dairy product including ice cream, soups, beverages, tea, drinks, alcoholic drinks and vitamin complex, etc. and other ordinary health food.

A composition for health promoting beverages can additionally include various flavors or natural carbohydrates, like any other ordinary beverages. Natural carbohydrates are exemplified by monosaccharides such as glucose and fructose, disacchsrides such as maltose and sucrose, polysaccharides such as dextrin, cyclodextrin, and sugar alcohols such as xilytole, sorbitol and

erythritol. As a sweetening agent, natural sweeteners such as thaumatin and stevia extract, and synthetic sweeteners such as saccharin and aspartame can be used. It is preferred to add natural carbohydrates by 0.1 - 20 g per 100 ml of a composition of the present invention, and is more preferred to add 1 - 10 g of natural carbohydrates to 100 ml of the composition.

In addition to the above, a composition of the present invention can also include various nutrients, vitamins, electrolytes, flavoring agents, coloring agents, pectic acid and its salts, alginic acid and its salts, organic acids, protective colloidal thickeners, pH regulators, stabilizers, antiseptics, glycerin, alcohol, carbonating agents used in carbonated beverages, etc. The composition of the present invention can further include sarcocarps to produce natural fruit juices, fruit beverages and vegetable beverages. Each ingredient is used either independently or in combination with others. At this time, the mixing rate is not so important but in general, 0.05 - 50 parts of weight per 100 parts of weight of the composition of the present invention is preferred.

EXAMPLES

Practical and presently preferred embodiments of the present invention are illustrative as shown in the following Examples.

However, it will be appreciated that those skilled in the art, on consideration of this disclosure, may make modifications and improvements within the spirit and scope of the present invention.

Example 1: Preparation of *Notoginseng radix* extract

Cultivated *Notoginseng radix* was purchased from a wholesale dried medicinal herb store.

<1-1> Preparation of *Notoginseng radix* crude extract

<1-1-1> Crude alcohol extract of *Notoginseng radix*

Notoginseng radix was cut into 1~2 cm fragments. The fragments were washed with running water to eliminate impurities. The fragments were pulverized. 200 g. of the *Notoginseng radix* powder was put in a 3 l flask, which was stirred at reflux at 78.5°C using 2,000 ml of ethanol. Extraction by heating was repeated three times for 4 hours. The extract was filtered and vacuum-concentrated under reduced pressure by using vacuum rotary evaporator under 40°C, resulting in

Notoginseng radix crude extract containing 2.7 g of
Notoginseng radix powder (RF1M) (yield : 1.35%).

<1-1-2> Crude water extract of *Notoginseng radix*

5 *Notoginseng radix* crude extract was extracted by
the same method as described in the above <1-1-1> and
the only difference in the procedure was that water was
used instead of ethanol as an extraction solvent.

10 <1-1-3> Crude mixed solvent extract of *Notoginseng*
radix

Notoginseng radix crude extract was extracted by
the same method as described in the above <1-1-1> and
the only difference in the procedure was that a mixed
15 solvent of water (25%) and ethanol (75%) was used
instead of ethanol as an extraction solvent.

<1-2> Separation of *Notoginseng radix* crude extract

 A fraction (RF1MB) was obtained from the crude
20 extract (RF1M) prepared in the above <1-1-1> at room
temperature by using 500 ml of normal butanol (n-
butanol) as a solvent, for which a fraction funnel was
used and solvent fractionation was repeated three times.

RF1MB4 fraction was separated from the RF1MB .
fraction by column chromatography. Column
chromatography was performed again with the RF1MB4
fraction, resulting in the final fraction of
5 *Notoginseng radix* extract (RF1MB4b).

Extraction and separation method of *Notoginseng*
radix extract of the present invention is described in
FIG. 1.

10 In experimental examples of the invention, the
final extraction of *Notoginseng radix* extract (RF1MB4b)
was concentrated and then freeze-dried. The dried
fraction was diluted with water and used for *in vitro*
and animal tests.

15

Experimental Example 1: Inhibition of the release of
TNF- α by *Notoginseng radix* extract of the present
invention

20 Following experiments were performed to
investigate whether or not *Notoginseng radix* extract of
the present invention inhibited the release of TNF- α , a
cytokine separated from human monocytic cell line 'THP-
1 cell'.

<1-1> Cell selection and culture

The below cell line was used to investigate the effect of *Notoginseng radix* extract of the present invention on the release of TNF- α .

- 5 Human originated cell line THP-1 (ATCC No. TIB-202) was purchased from ATCC (Rockville, USA) and cultured in RPMI 1640 (Gibco, BRL, USA) medium supplemented with 10% FBS (fetal bovine serum).

10 <1-2> Quantification of released TNF- α

In order to investigate the effect of *Notoginseng radix* extract of the present invention on the release of TNF- α , the amount of released TNF- α was measured by ELISA using cells prepared in the above <1-1>.

- 15 Cells were plated into a 96-well plate by 5×10^5 cells/ml and lipopolysaccharide (LPS) was added in order to activate cells for the release of TNF- α .

An experimental group was treated with *Notoginseng radix* extract (RF1MB4b) at the
20 concentration of 2 or 10 $\mu\text{l/ml}$ together with LPS. After the treatment, the released TNF- α in culture supernatant was quantified by ELISA.

The results are presented in FIG. 2.

As shown in FIG. 2, when an experimental group

was treated with low concentration (2 $\mu\text{l/ml}$) of
Notoginseng radix extract (RF1MB4b), the amount of
released TNF- α of the experimental group was just a
little different from that of a control group not
5 treated with the extract. But, when the extract was
provided with high concentration (10 $\mu\text{l/ml}$), the amount
of released TNF- α in the experimental group was greatly
decreased, comparing to a control group.

Thus, the above results indicate that *Notoginseng*
10 *radix* extract of the present invention inhibits the
release of TNF- α .

Experimental Example 2: Selective apoptosis of
activated T-cells by *Notoginseng radix* extract of the
15 present invention

In order to confirm whether or not *Notoginseng*
radix extract of the present invention could destroy
activated T-cells only, following experiments were
performed.

20

<2-1> Separation and activation of T-cells

A lymph node of a 5-week-old female mouse was
taken out and mashed by the back tip of a sterilized

syringe to extract cells. The cells were filtered by a cell-filter (Falcon, NJ USA) and washed with PBS, then put in a culture medium at the concentration of 2×10^6 cells/ml. As a culture medium, RPMI 1640 (Gibco, BRL, USA) supplemented with 10% FBS (fetal bovine serum) was used.

In order to activate T-cells only, concanavalin A was added by $5 \mu\text{g/ml}$ to the medium, followed by culture for 48 hours. After 48 hours of culture, 10 mg/ml of methyl- α -D-mannopyranoside (sigma, Germany) was put in the medium, followed by further culture for 30 minutes. Then, the cells were washed with PBS three times and put in a culture medium supplemented with 100 units/ml of human interleukine-2 (hIL-2, R&D, MN, USA), followed by further culture for 48 hours and cell density was maintained as 2×10^6 cells/ml during the culture (Lenardo MJ. et al. : Interleukin-2 programs mouse alpha beta T lymphocytes for apoptosis. Nature. 353(6347):858-61. 1991).

20

<2-2> Investigation of selective apoptosis of activated T-cells

The concentration of activated T-cells was adjusted to 1×10^6 cells/ml, then they were put in a 96-

well plate (Falcon, USA) by 200 μ l/well. At that time, 100 units/ml of human interleukine-2 (hIL-2) was added to each well.

While a control group was not treated with
5 *Notoginseng radix* extract, an experimental group was treated with the final fraction (RF1MB4b) of *Notoginseng radix* extract prepared in the above example at different concentrations (5 μ g/ml, 10 μ g/ml, 20 μ g/ml) before being cultured for 24 hours.

10 As a control, inactivated cells were prepared as follows.

Single cells were collected from spleen and cell density was adjusted to 2×10^6 cells/ml, which were distributed to a 96 well plate by 200 μ l/well.
15 *Notoginseng radix* extract was added thereto, followed by culture for 24 hours. After 24 hours of culture, the cells were transferred to a flow tube, to which propidium iodide (PI) was added. Then, live cells were counted for 20 seconds by using CellQuest program of
20 FACSCaliver (Becton Dickinson, France).

Apoptosis was calculated as follows : $(1 - F \text{ extract treated cells} / \text{untreated cells}) \times 100$. All candidate drugs were examined by that math formula to choose a drug to induce high apoptosis of activated T-
25 cells but low apoptosis of naive T-cells (Sabapathy K,

Hu Y, Kallunki T, Schreiber M, David JP, Jochum W, Wagner EF, Karin M. : JNK2 is required for efficient T-cell activation and apoptosis but not for normal lymphocyte development. Curr Biol. 11;9(3):116-25.

5 1999).

The results are presented in FIG. 3.

As shown in FIG. 3, when *Notoginseng radix* extract of the present invention was treated with high concentration over 5 $\mu\text{l/ml}$, activated T-cells were selectively destroyed while inactivated T-cells still remained.

10

Thus, it was confirmed that *Notoginseng radix* extract of the present invention destroys activated T-cells selectively and the apoptosis effect was concentration-dependent.

15

Experimental Example 3: Inhibition of the progress of arthritis in test animals with type 2 collagen induced arthritis by *Notoginseng radix* extract of the present invention

20

In order to investigate whether or not *Notoginseng radix* extract of the present invention could inhibit the progress of arthritis in test animals

having type 2 collagen induced arthritis, following experiments were performed.

<3-1> Inducement of arthritis in test animals

5 In order to prepare test animals having type 2 collagen induced arthritis, 5-6 week old male DBA1 mice were purchased from SCI company, Japan, and the mice were raised at 21°C with 40% humidity.

10 Bovine type 2 collagen (Condrex Co., Japan) was dissolved in 0.05% acetic acid, making the concentration 2 mg/ml. Then the type 2 collagen was mixed with the same amount of complete adjuvant (Condrex Co., Japan). While cooling down with ice, the
15 mixture became homogeneous suspension by using T-connector linked to 3 ml syringe. After confirming the suspension was prepared rightly, tail head of a mouse was sterilized with alcohol cotton and 100 µl of collagen suspension was injected under the skin of the
20 tail head.

<3-2> Oral administration of *Notoginseng radix* extract

(RF1MB4b) of the present invention

Notoginseng radix extract (RF1MB4b) prepared in

the above example was dissolved in water, resulting in 2.5 mg/ml solution. The solution was filtered by 0.25 uM filter.

5 The filtered solution was diluted to 0.2 mg/ml and was administered to the mouth of a mouse through sonde linked to a 1 ml syringe, once a day and by 0.05 mg/250 µl/mouse.

<3-3> Progress of arthritis : Naked eye observation and

10 diagnosis

In order to investigate arthritis treating effect of *Notoginseng radix* extract (RF1MB4b) of the present invention, the *Notoginseng radix* extract (RF1MB4b) prepared in the above example was administered by the
15 same method as described in the above <3-2> to test animals having arthritis induced by the injection of collagen suspension.

Arthritis was developed 30 days after collagen suspension was injected to a mouse. Naked eye
20 observation on lesion of arthritis was performed by using following scores based on literature cited.

0 : No swelling or flair, 1 : Light swelling and flair in joint, 2: Clear swelling and flair in joint,

3: Severe swelling and flair in joint including knuckle joint, 4 : Severe swelling in all over the joint.

Therefore, the highest score of lesion of arthritis is 16 per mouse, which sums up scores of forelegs and hind legs, and the highest score per one leg is 4 (Courtenay JS, Dallman MJ, Dayan AD, et al. : Immunization against heterologous type II collagen induces arthritis in mice. Nature 283: 666-668. 1980).

FIG. 4 and FIG. 5 present the results of investigation, after oral administration of the extract, of arthritis progress inhibiting effect of *Notoginseng radix* extract of the present invention in test animals with type 2 collagen induced arthritis.

In FIG. 4, the arthritis progress inhibiting effect of *Notoginseng radix* extract of the present invention in test animals with type 2 collagen induced arthritis was presented as arthritis index, and FIG. 5 is a set of photographs showing the arthritis progress inhibiting effect of *Notoginseng radix* extract of the present invention in animals having type 2 collagen induced arthritis.

As shown in FIG. 4, when *Notoginseng radix* extract of the present invention was orally administered into a mouse having type 2 collagen induced arthritis, the progress of the disease was

obviously inhibited from the 9th day of administration, comparing to a control group.

As shown in FIG. 5, both a control medicine without *Notoginseng radix* extract and an experimental
5 medicine including the extract were orally administered respectively to mice having type 2 collagen induced arthritis. Big difference between the two was observed after 21 days from the administration. A mouse treated with a control medicine showed very severe swelling all
10 over the joints but a mouse administered with an experimental medicine just showed light flair and swelling in joints.

Therefore, it was confirmed that *Notoginseng radix* extract of the present invention effectively
15 inhibits the progress of arthritis.

Example 4: Acute toxicity test with *Notoginseng radix*
extract of the present invention

Notoginseng radix extract of the present
20 invention is classified into a food material, indicating that it is safe. But, for the use as a treatment medicine, acute toxicity of the extract had to be investigated as follows.

6-week old SPF mice were used in the tests for acute toxicity. *Notoginseng radix* extract (RF1MB4b) prepared in the above example was suspended in distilled water and orally administered once to 5 mice per group at the dosage of 2, 1, and 0.5 g/kg.

Death, clinical symptoms, and weight change in mice were observed, hematological tests and biochemical tests of blood were performed, and any abnormal signs in the gastrointestinal organs of chest and abdomen were checked with eyes during autopsy.

The results showed that *Notoginseng radix* extract of the present invention did not cause any specific clinical symptoms, weight change, or death in mice. No change was observed in hematological tests, biochemical tests of blood, and autopsy.

Notoginseng radix extract (RF1MB4b) of the present invention used in this experiment is evaluated to be safe substance since it does not cause any toxic change in mice up to the level of 2 g/kg and its estimated LD₅₀ values are much greater than 2 g/kg in mice.

Manufacturing Example 1: Preparation of pharmaceutical
formulations

<1-1> Preparation of powders

	<i>Notoginseng radix</i> extract	2g
5	Lactose	1g

Powders were prepared by mixing all the above components and filled airtight bag with them.

<1-2> preparation of tablets

10	<i>Notoginseng radix</i> extract	100 mg
	Corn starch	100 mg
	Lactose	100 mg
	Magnesium stearate	2 mg

Tablets were prepared by mixing all the above
15 components by the conventional method for preparing tablets.

<1-3> Preparation of capsules

	<i>Notoginseng radix</i> extract	100 mg
20	Corn starch	100 mg
	Lactose	100 mg
	Magnesium stearate	2 mg

Capsules were prepared by mixing the components above and filled gelatin capsules with them according to the conventional method for capsules.

5 Manufacturing Example 2: Preparation of food

Foodstuff containing *Notoginseng radix* extract of the present invention was prepared as follows.

<2-1> Preparation of cooking spices

10 Health improving spices and condiments containing *Notoginseng radix* extract of the present invention by 20-95 weight% were prepared.

<2-2> Preparation of tomato ketchup and sauce

15 Health improving tomato ketchup or sauce was prepared by adding *Notoginseng radix* extract of the present invention by 0.2-1.0 weight% to original tomato ketchup or sauce.

20 <2-3> Preparation of flour food

Health improving flour food was prepared by adding *Notoginseng radix* extract of the present invention by 0.5-5.0 weight% to wheat flour and then

making the flour into bread, cakes, cookies, crackers and noodles.

<2-4> Preparation of soups and gravies

5 *Notoginseng radix* extract of the present invention was added by 0.1-5.0 weight% to soups and gravies to produce health improving processed meats, noodle soups and gravies.

10 <2-5> Preparation of ground beef

Notoginseng radix extract of the present invention was added by 10 weight% to ground beef to prepare health improving ground beef.

15 <2-6> Preparation of dairy products

Notoginseng radix extract of the present invention was added by 5-10 weight% to milk to prepare health improving dairy products such as butter, ice cream, etc.

20

<2-7> Preparation of Sunsik

 Brown rice, barley, glutinous rice and coix (job's tear) were gelatinized by the conventional

method, followed by drying. The dried mixture was distributed and pulverized, resulting in 60-mesh grain size of powders.

5 Black bean, black sesame and perilla were steamed and dried by the conventional method. The dried mixture was distributed and pulverized, resulting in 60-mesh grain size of powders.

10 *Notoginseng radix* extract of the present invention was vacuum-concentrated under reduced pressure using a vacuum concentrator, which was then spray-dried with a hot-air drier. The dried material was pulverized by a grinder, resulting in 60-mesh grain size of powders.

15 The prepared grain, seeds, and dried *Notoginseng radix* extract powders were all mixed at the following ratio.

Grain (brown rice 30 weight%, coix 15 weight%, barley 20 weight%),

20 Seeds (perilla 7 weight%, black bean 8 weight%, black sesame 7 weight%),

Dried powder of *Notoginseng radix* extract (3 weight%),

Ganoderma lucidum (0.5 weight%),

25 *Rehmannia glutinosa* (0.5 weight%)

Manufacturing Example 3: Preparation of beverages

<1-1> Preparation of carbonated beverages

Sugar (5-10%), citric acid (0.05-0.3%), caramel
5 (0.005-0.02%) and vitamin C (0.1-1%) were mixed, to
which purified water (79-94%) was added to make syrup.
The prepared syrup was sterilized at 85-98°C for 20-180
seconds, then mixed with cooling water at the ratio of
1 : 4. Then, carbon dioxide gas (0.5-0.82%) was given
10 to the mixture to prepare carbonated beverages
containing *Notoginseng radix* extract of the present
invention.

<1-2> Preparation of health beverages

15 Acid fructose (0.5%), oligosaccharide (2%), sugar
(2%), salt (0.5%) and water (75%) were all mixed with
Notoginseng radix extract evenly, followed by
sterilization. The mixture was put in a small
container such as a glass bottle or pat bottle,
20 resulting in health beverages.

<1-3> Preparation of vegetable juice

5 g of *Notoginseng radix* extract of the present invention was added to 1,000 ml of tomato or carrot juice to prepare health vegetable juice.

5 <1-4> Preparation of fruit juice

1 g of *Notoginseng radix* extract of the present invention was added to 1,000 ml of apple or grape juice to produce health fruit juice.

10

INDUSTRIAL APPLICABILITY

As explained hereinbefore, *Notoginseng radix* extract of the present invention has activities of inhibiting TNF- α release and destroying activated T-cells selectively.

15

Therefore, *Notoginseng radix* extract of the present invention can be effectively used for the production of health food or a medicine for preventing and treating arthritis.

What is claimed is

1. A composition for preventing and treating
arthritis comprising a purified form of butanol
fraction of ethanol extract of *Notoginseng radix*
as an effective ingredient.
2. A composition as claimed in claim 1, the purified
form of butanol fraction of ethanol extract of
Notoginseng radix is produced by the steps of:
obtaining ethanol extract of *Notoginseng radix* by
using ethanol as an extraction solvent; obtaining
butanol fraction of ethanol extract from the
ethanol extract by using butanol as fractionation
solvent; separating an intermediate form of
butanol fraction of ethanol extract from the
butanol fraction by using column chromatography;
and separating a purified form of butanol fraction
of ethanol extract from the intermediate form of
butanol fraction by using column chromatography.
3. A pharmaceutical composition for preventing and
treating arthritis comprising the composition of
any one of claim 1 to claim 2.

4. A health food composition for preventing and treating arthritis comprising the composition of any one of claim 1 to claim 2.

ABSTRACT OF THE DISCLOSURE

The present invention relates to a composition comprising *Notoginseng radix* extract for preventing and treating arthritis as an effective ingredient.

5 *Notoginseng radix* extract of the present invention inhibits release of tumor necrosis factor-alpha (TNF- α) and is the death of activated T-cells only, so that it can be effectively used for the production of a medicine for preventing and treating
10 arthritis and health food as well.